

Second Semester B.E. Degree Examination, December 2011
Engineering Mathematics - II

Time: 3 hrs.

Max. Marks: 100

Note: 1. Answer any FIVE full questions, choosing at least two from each part.

- 2. Answer all objective type questions only on OMR sheet page 5 of the answer booklet.**
3. Answer to objective type questions on sheets other than OMR will not be valued.

PART - A

- 1** a. Choose your answers for the following : (04 Marks)
- The general solution of the equation $yp^2 + (x-y)p - x = 0$ is

A) $(x-y-c)(x^2 + y^2 - c) = 0$	B) $(y-x-c)(x^2 - y^2 - c) = 0$
C) $(y-x-c)(y^2 - x^2 - c) = 0$	D) $(y-x-c)(x^2 + y^2 - c) = 0$
 - The given differential equation is solvable for x, if it is possible to express x in terms of,

A) x and y	B) x and p	C) y and p	D) None of these
------------	------------	------------	------------------
 - The singular solution of the equation $y = px + \frac{a}{p}$ is

A) $y^2 = 4ax$	B) $x^2 = 4ay$	C) $x^2 = y$	D) $y^2 = x$
----------------	----------------	--------------	--------------
 - The general solution of Clairaut's equation is,

A) $y = cx + f(c)$	B) $x = cy + f(c)$	C) $y = cx - f(c)$	D) None of these
--------------------	--------------------	--------------------	------------------
- b. Solve : $p(p+y) = x(x+y)$. (04 Marks)
- c. Obtain the general solution and the singular solution of the equation, $y = 2px + p^2y$. (06 Marks)
- d. Obtain the general and singular solution of Clairaut's equation, $xp^3 - yp^2 + 1 = 0$. (06 Marks)
- 2** a. Choose your answers for the following : (04 Marks)
- The particular integral of $(D^2 + a^2)y = \sin ax$ is

A) $-\frac{x}{2a} \cos ax$	B) $\frac{x}{2a} \cos ax$	C) $-\frac{ax}{2} \cos ax$	D) $\frac{ax}{2} \cos ax$
----------------------------	---------------------------	----------------------------	---------------------------
 - The solution of the differential equation $y'' + y = 0$ satisfying the conditions $y(0) = 1$ and $y\left(\frac{\pi}{2}\right) = 2$ is

A) $y = \cos x - 2 \sin x$	B) $y = 2 \sin x - \cos x$
C) $y = \cos x + 2 \sin x$	D) $y = C_1 \cos x + C_2 \sin x$
 - P.I of $(D+1)^2y = xe^{-x}$ is,

A) $\frac{x}{6}e^{-x}$	B) $\frac{x^3}{6}e^{-x}$	C) $-\frac{x^3}{6}e^{-x}$	D) $\frac{x^2}{2}e^{-x}$
------------------------	--------------------------	---------------------------	--------------------------
 - P.I of $(D^2 + D)y = x^2 + 2x + 4$ is

A) $\frac{x^2}{3} + 4x$	B) $\frac{x^3}{3} + 4$	C) $\frac{x^3}{3} + 4x$	D) $\frac{x^3}{3} + 4x^2$
-------------------------	------------------------	-------------------------	---------------------------
- b. Solve : $(D-2)^2y = 8(e^{2x} + \sin 2x)$ (04 Marks)
- c. Solve : $y'' - 2y' + y = x \cos x$ (06 Marks)
- d. Solve : $\frac{dx}{dt} - 7x + y = 0$, $\frac{dy}{dt} - 2x - 5y = 0$. (06 Marks)

3 a. Choose your answers for the following : (04 Marks)

- The complementary function of the equation $x^2y'' - xy' + y = \log x$ is
 A) $y = (C_1 + C_2x)e^x$
 B) $y = (C_1 + C_2 \log x)x$
 C) $y = (C_1 + C_2x)x$
 D) $y = C_1e^x + C_2e^{-x}$
- The homogeneous linear differential equation whose auxillary equation has roots 1, -1 is
 A) $x^2y_2 - xy_1 + y = 0$
 B) $x^2y_2 - xy_1 - y = 0$
 C) $y'' - y = 0$
 D) $x^2y_2 + xy_1 - y = 0$
- To transform $xy'' + y' = \frac{1}{x}$ into a linear differential equation with constant coefficients put $x = \dots\dots$
 A) e^t
 B) e^{-t}
 C) $\log t$
 D) None of these
- The solution of $x^2y'' + xy' = 0$ is
 A) $y = C_1 \cos x + C_2 \sin x$
 B) $y = C_1 e^x + C_2 e^{-x}$
 C) $y = a \log x + b$
 D) $y = C_1 + 6x^3$

b. Solve $y'' - 6y' + 9y = \frac{e^{3x}}{x^2}$ by the method of variation of parameters. (04 Marks)

c. Solve : $(1+x^2)\frac{d^2y}{dx^2} + (1+x)\frac{dy}{dx} + y = 2\sin[\log(1+x)]$. (06 Marks)

d. Solve by Frobenius method the equation: $4x\frac{d^2y}{dx^2} + 2\frac{dy}{dx} + y = 0$. (06 Marks)

4 a. Choose your answers for the following : (04 Marks)

- The solution of $\frac{\partial^2 z}{\partial y^2} = \sin(xy)$ is
 A) $z = -x^2 \sin(xy) + yf(x) + g(x)$
 B) $z = -x^2 \cos(xy) - yf(x) + g(x)$
 C) $z = -\frac{\sin(xy)}{x^2} + yf(x) + g(x)$
 D) None of these
- A solution of $(y-z)p + (z-x)q = x-y$ is
 A) $x^2 + y^2 + z^2 = f(x-y-z)$
 B) $x^2 + y^2 + z^2 = f(x+y+z)$
 C) $x^2 - y^2 - z^2 = f(x+y+z)$
 D) $x^2 + y^2 - z^2 = f(x+y+z)$
- The partial differential equation obtained from $z = ax + by + ab$ is
 A) $px + qy + z = 0$
 B) $px + qy + z^2 = 0$
 C) $px - qy = z$
 D) $px + qy = z$
- The partial differential equation obtained from $z = e^y f(x+y)$ is
 A) $p + z = q$
 B) $p - z = q$
 C) $p - q = z$
 D) None of these

b. Form the partial differential equation by eliminating the arbitrary functions from $z = f(y-2x) + g(2y-x)$. (04 Marks)

c. Solve : $(x^2 - yz)p + (y^2 - zx)q = z^2 - xy$. (06 Marks)

d. Solve : $4\frac{\partial u}{\partial x} + \frac{\partial u}{\partial y} = 3u$ by the method of separation of variables, given $u(0, y) = 2e^{5y}$. (06 Marks)

PART - B

5 a. Choose your answers for the following : (04 Marks)

- $\int\limits_0^2 \int\limits_0^x (x+y) dx dy = \dots\dots$
 A) 0
 B) 1
 C) 3
 D) 4

ii) $\int_0^{\infty} e^{-x^2} dx = \dots$

- A) $\sqrt{\pi}$ B) $\frac{\sqrt{\pi}}{2}$ C) $\sqrt{\frac{\pi}{2}}$ D) $\frac{\pi}{2}$

iii) The value of $\beta(2, 1) + \beta(1, 2)$ is

- A) 0 B) $\frac{1}{2}$ C) 2 D) 1

iv) $\iiint_{0 \ 1 \ 1}^{2 \ 3 \ 2} xy^2 z \ dz \ dy \ dx = \dots$

- A) 26 B) 25 C) 1 D) 0

b. Change the order of integration in $\int_0^1 \int_{x^2}^{2-x} xy \ dx \ dy$ and hence evaluate the same. (04 Marks)

c. Evaluate $\int_0^{\infty} \int_0^{\infty} e^{-(x^2+y^2)} dx \ dy$ by changing to polar coordinates. (06 Marks)

d. Show that $\beta(m, n) = \int_0^1 \frac{x^{m-1} + x^{n-1}}{(1+x)^{m+n}} dx$. (06 Marks)

6. a. Choose your answers for the following : (04 Marks)

i) If $\vec{F} = x^2 \mathbf{i} + xy \mathbf{j}$ then $\int_C \vec{F} \cdot d\vec{r}$ from $(0, 0)$ to $(1, 1)$ along the line $y = x$ is

- A) 0 B) $\frac{2}{3}$ C) $\frac{3}{2}$ D) None of these

ii) The value of $\iint_S (yz \ dy \ dz + zx \ dz \ dx + xy \ dx \ dy)$ where S is the surface of unit sphere $x^2 + y^2 + z^2 = 1$ is

- A) 0 B) 4π C) $\frac{4\pi}{3}$ D) 10π

iii) A necessary and sufficient condition that the line integral $\int_L \vec{F} \cdot d\vec{R}$ for every closed curve C is

- A) $\text{Curl } \vec{F} = 0$ B) $\text{div } \vec{F} = 0$ C) $\text{Curl } \vec{F} \neq 0$ D) $\text{div } \vec{F} \neq 0$

iv) If V is the volume bounded by a surface S and \vec{F} is a continuously differentiable vector function then $\iiint_V \text{div } \vec{F} \ dv = \dots$

- A) 0 B) $\iint_S \vec{F} \cdot \hat{n} \ ds$ C) $\iint_S \vec{F} \cdot \hat{n} \ ds$ D) None of these

b. Using Green's theorem evaluate $\int_C [(xy + y^2) dx + x^2 dy]$ where C is bounded by $y = x$ and $y = x^2$. (04 Marks)

c. Verify Stoke's theorem for the vector $\vec{F} = (x^2 + y^2) \mathbf{i} - 2xy \mathbf{j}$ taken round the rectangle bounded by $x = 0, x = a, y = 0, y = b$. (06 Marks)

d. Using divergence theorem evaluate $\int_S \vec{F} \cdot d\vec{S}$ where $\vec{F} = 4xi - 2y^2j + z^2k$ and S is the surface bounded by the region $x^2 + y^2 = 4, z = 0, z = 3$. (06 Marks)

- 7 a. Choose your answers for the following : (04 Marks)
- If $L\{f(t)\} = f(s)$ then $L\{e^{-at}f(t)\}$ is
 A) $f(s-a)$ B) $f(s+a)$ C) $f(s)$ D) None of these
 - $L\left\{\frac{\sin at}{t}\right\} = \dots$
 A) $\cos^{-1}\left(\frac{s}{a}\right)$ B) $\tan^{-1}\frac{s}{a}$ C) $\frac{\pi}{2} + \tan^{-1}\frac{s}{a}$ D) None of these
 - $L\{u(t+2)\} = \dots$
 A) $\frac{e^{-2s}}{s^2}$ B) e^{2s} C) $\frac{e^{2s}}{s}$ D) $\frac{e^{-2s}}{s}$
 - $L\{s(t)\} = \dots$
 A) 0 B) e^{-as} C) ∞ D) 1
- b. Find the value of $\int_0^\infty t^3 e^{-t} \sin t dt$ using Laplace transforms. (04 Marks)
- c. If $f(t) = \begin{cases} t, & 0 \leq t \leq a \\ 2a-t, & a \leq t \leq 2a \end{cases}$, where $f(t+2a) = f(t)$, show that $L\{f(t)\} = \frac{1}{s^2} \tan h\left(\frac{as}{2}\right)$. (06 Marks)
- d. Express $f(t) = \begin{cases} 1, & 0 < t \leq 1 \\ t, & 1 < t \leq 2 \\ t^2, & t > 2 \end{cases}$ in terms of unit step function and hence find its Laplace transform. (06 Marks)
- 8 a. Choose your answers for the following : (04 Marks)
- $L^{-1}\left\{\frac{1}{s^n}\right\}$ is possible only when n is
 A) zero B) -ve integer C) +ve integer D) -ve rational
 - $L^{-1}\left\{\frac{s}{(s-1)^3}\right\} = \dots$
 A) $e^{-t}(t+t^2)$ B) $e^t\left(t+\frac{t^2}{2!}\right)$ C) $t e^t + t^2 e^t$ D) None of these
 - $L^{-1}\left\{\log\left(\frac{s+1}{s-1}\right)\right\} = \dots$
 A) $2 \sin t$ B) $2 \cos h t$ C) $\sin h t$ D) $2 \sin h t$
 - $L^{-1}\left\{\frac{s}{(2s+3)^2}\right\} = \dots$
 A) $-\frac{1}{8}(2-3t)e^{\frac{-3t}{2}}$ B) $\frac{1}{8}(2-3t)e^{\frac{-3t}{2}}$ C) $2e^{\frac{-3t}{2}} - 3te^{\frac{-3t}{2}}$ D) None of these
- b. Find $L^{-1}\left\{\frac{5s+3}{(s-1)(s^2+2s+5)}\right\}$. (04 Marks)
- c. Using convolution theorem evaluate $L^{-1}\left\{\frac{s^2}{(s^2+a^2)(s^2+b^2)}\right\}$. (06 Marks)
- d. Solve $y''' + 2y'' - y' - 2y = 0$ given $y(0) = y'(0) = 0$ and $y''(0) = 6$ by using Laplace transform method. (06 Marks)

* * * * *